

e/p

N64-24965

Global Distribution of the Net Energy Balance of the
Atmosphere from Tiros Radiation Data

Code None

S. I. Rasool

Reports

Global Distribution of the Net Energy Balance of the Atmosphere from Tiros Radiation Data

Abstract. *The radiation data from Tiros II and Tiros III have been analyzed to obtain the monthly averages of the global distribution of the total outgoing radiation from the earth. These data have been combined with climatological estimates of the incoming radiation for the same periods to obtain monthly averages of the regional distribution of the net energy balance. The results show that these data could be useful in studies of the meridional circulation and the evolution of large-scale weather systems.*

We have carried out an analysis of the radiation data acquired by the Tiros meteorological satellites, in order to determine the regional and time variations in the energy balance of the earth and its atmosphere.

The energy balance of the earth-atmosphere system is made up of the difference between the incoming solar radiation, mostly in the visible, and the outgoing terrestrial radiation in the infrared. It is well known that the latitudinal variation of the energy balance shows an excess of incoming solar radiation over outgoing radiation near the equator, and a deficiency at the poles. It is this variation of the energy balance with latitude that drives the atmospheric heat engine. Thus, through the determination of the latitudinal average of the energy balance with the aid of the Tiros data, we obtain the information which is necessary to understand the general circulation of the atmosphere.

At the same time, we have obtained some information regarding the regional variations in the energy balance of the earth-atmosphere system. Knowledge of these variations gives us a very important source of information for the understanding of large-scale weather systems.

Simpson (1) carried out the first extensive studies of the atmospheric energy balance, by the application of a simplified radiation theory to empirical physical models of the atmosphere. This problem has since been re-examined by Houghton (2), London (3), Budyko (4), and others.

The radiation instrumentation in Tiros, and the physical significance of this experiment, have already been discussed in detail by Bandeen *et al.* (5) and Nordberg *et al.* (6).

Three of the five channels of the Tiros radiometer measure terrestrial radiation in the far infrared, corresponding to wavelength intervals of 5.8 to 6.5 μ , 7 to 13 μ , and 7 to 30 μ . The other two channels record the solar radiation reflected by the earth, in the visible spectrum, which gives an estimate of the albedo of any region.

In Tiros II and Tiros III, which were active for 5 and 3 months, respectively, the channels which measure the reflected visible energy degraded very quickly, and almost no data are available for these channels. It is also possible to obtain information regarding the solar input by the statistical analysis of the Tiros cloud-cover photographs (7), but this analysis is still in a preliminary stage. Therefore, for this first study of the energy balance, we have used the climatological estimates of Budyko (4) for the incoming radiation. These give the monthly averages of solar radiation reaching the ground as a function of latitude and longitude, as derived from ground-based pyroheliometer data.

For the outgoing radiation we have used the Tiros data in the infrared channels. Channel 4, which is sensitive in the 7- to 30- μ interval, and thus records almost 80 percent of the total outgoing radiation from the earth, also degraded after approximately 1 month in each of the satellites. The data from channel 2, sensitive in the 7- to 13- μ

interval, however, is available from 23 November 1960 to 12 April 1961 for Tiros II and from 12 July to 10 September 1961 for Tiros III. There was some degradation in this case also, but fairly good estimates of the correction factors for this channel are now available.

Wark *et al.* (8) have shown that data from this channel can be used to obtain reasonably good estimates of the total outgoing radiation from the earth. Numerical factors for converting channel 2 measurements to total outgoing radiation are now available for both Tiros II and III. A comparison between the values of total outgoing radiation derived from channels 4 and 2 for the overlapping period give satisfactory results.

To obtain an estimate of the temporal as well as regional variations in the total outgoing flux, we have divided the surface of the earth into intervals of 10° latitude by 10° longitude between 50°N and 50°S. Because the inclination of the satellite orbit is 48° the data for polar regions are not available. All measurements made by the satellite in each of these intervals, at nadir angles less than 25°, have been converted to the total outgoing flux and averaged separately for each month. There are approximately 500 observation points per month in each 10° square.

The total outgoing flux estimated in this manner is, however, liable to have a small diurnal bias. This is because the orbit of the satellite precesses in right ascension at a rate of 6° per day, or it has a precession period of 9 weeks. Therefore our results for the total outgoing radiation are affected by the diurnal variation of the ground temperature. However, this effect is small for the particular case of this study which is concerned with the long-period averages over large areas. Because of the cloudiness and the presence of water vapor, the average of the channel-2 data over large regions and extended periods of time does not give the ground temperature, but has been shown (9) to give the effective temperature at a height of about 3 km, and therefore the effect of the diurnal variation is much reduced.

As mentioned earlier, for the incoming radiation we have used the climatological estimates of Budyko (4), which are based on the ground observations of solar radiation, and are given

as monthly averages, also in 10° by 10° grids. These values were corrected for the absorption of solar radiation by the atmosphere, which, according to London (3), varies with latitude and

season and is about 30 to 40 percent of the radiation which reaches the ground. The energy balance for each 10° by 10° grid, that is, the difference between these corrected values of in-

coming energy and the Tiros-measured values of the outgoing energy, is plotted on a month by month basis in Figs. 1 through 6.

These correspond to the months of




Fig. 1. December 1960.




Fig. 2. January 1961.




Fig. 3. February 1961.




Fig. 4. March 1961.




Fig. 5. July 1961.




Fig. 6. August 1961.

Figs. 1-6. Distribution of net energy balances for the months of December 1960 (Fig. 1), January 1961 (Fig. 2), February 1961 (Fig. 3), March 1961 (Fig. 4), July 1961 (Fig. 5), and August 1961 (Fig. 6). Extreme dark shade $\sim > +1.75 \times 10^7$ ergs $\text{cm}^{-2} \text{sec}^{-1}$ and extreme light shade $< -1.65 \times 10^7$ ergs $\text{cm}^{-2} \text{sec}^{-1}$.

Reports

Global Distribution of the Net Energy Balance of the Atmosphere from Tiros Radiation Data

Abstract. *The radiation data from Tiros II and Tiros III have been analyzed to obtain the monthly averages of the global distribution of the total outgoing radiation from the earth. These data have been combined with climatological estimates of the incoming radiation for the same periods to obtain monthly averages of the regional distribution of the net energy balance. The results show that these data could be useful in studies of the meridional circulation and the evolution of large-scale weather systems.*

We have carried out an analysis of the radiation data acquired by the Tiros meteorological satellites, in order to determine the regional and time variations in the energy balance of the earth and its atmosphere.

The energy balance of the earth-atmosphere system is made up of the difference between the incoming solar radiation, mostly in the visible, and the outgoing terrestrial radiation in the infrared. It is well known that the latitudinal variation of the energy balance shows an excess of incoming solar radiation over outgoing radiation near the equator, and a deficiency at the poles. It is this variation of the energy balance with latitude that drives the atmospheric heat engine. Thus, through the determination of the latitudinal average of the energy balance with the aid of the Tiros data, we obtain the information which is necessary to understand the general circulation of the atmosphere.

At the same time, we have obtained some information regarding the regional variations in the energy balance of the earth-atmosphere system. Knowledge of these variations gives us a very important source of information for the understanding of large-scale weather systems.

Simpson (1) carried out the first extensive studies of the atmospheric energy balance, by the application of a simplified radiation theory to empirical physical models of the atmosphere. This problem has since been re-examined by Houghton (2), London (3), Budyko (4), and others.

The radiation instrumentation in Tiros, and the physical significance of this experiment, have already been discussed in detail by Bandeen *et al.* (5) and Nordberg *et al.* (6).

Three of the five channels of the Tiros radiometer measure terrestrial radiation in the far infrared, corresponding to wavelength intervals of 5.8 to 6.5 μ , 7 to 13 μ , and 7 to 30 μ . The other two channels record the solar radiation reflected by the earth, in the visible spectrum, which gives an estimate of the albedo of any region.

In Tiros II and Tiros III, which were active for 5 and 3 months, respectively, the channels which measure the reflected visible energy degraded very quickly, and almost no data are available for these channels. It is also possible to obtain information regarding the solar input by the statistical analysis of the Tiros cloud-cover photographs (7), but this analysis is still in a preliminary stage. Therefore, for this first study of the energy balance, we have used the climatological estimates of Budyko (4) for the incoming radiation. These give the monthly averages of solar radiation reaching the ground as a function of latitude and longitude, as derived from ground-based pyroheliometer data.

For the outgoing radiation we have used the Tiros data in the infrared channels. Channel 4, which is sensitive in the 7- to 30- μ interval, and thus records almost 80 percent of the total outgoing radiation from the earth, also degraded after approximately 1 month in each of the satellites. The data from channel 2, sensitive in the 7- to 13- μ

interval, however, is available from 23 November 1960 to 12 April 1961 for Tiros II and from 12 July to 10 September 1961 for Tiros III. There was some degradation in this case also, but fairly good estimates of the correction factors for this channel are now available.

Wark *et al.* (8) have shown that data from this channel can be used to obtain reasonably good estimates of the total outgoing radiation from the earth. Numerical factors for converting channel 2 measurements to total outgoing radiation are now available for both Tiros II and III. A comparison between the values of total outgoing radiation derived from channels 4 and 2 for the overlapping period give satisfactory results.

To obtain an estimate of the temporal as well as regional variations in the total outgoing flux, we have divided the surface of the earth into intervals of 10° latitude by 10° longitude between 50°N and 50°S. Because the inclination of the satellite orbit is 48° the data for polar regions are not available. All measurements made by the satellite in each of these intervals, at nadir angles less than 25°, have been converted to the total outgoing flux and averaged separately for each month. There are approximately 500 observation points per month in each 10° square.

The total outgoing flux estimated in this manner is, however, liable to have a small diurnal bias. This is because the orbit of the satellite precesses in right ascension at a rate of 6° per day, or it has a precession period of 9 weeks. Therefore our results for the total outgoing radiation are affected by the diurnal variation of the ground temperature. However, this effect is small for the particular case of this study which is concerned with the long-period averages over large areas. Because of the cloudiness and the presence of water vapor, the average of the channel-2 data over large regions and extended periods of time does not give the ground temperature, but has been shown (9) to give the effective temperature at a height of about 3 km, and therefore the effect of the diurnal variation is much reduced.

As mentioned earlier, for the incoming radiation we have used the climatological estimates of Budyko (4), which are based on the ground observations of solar radiation, and are given

December 1960 and January, February, March, July, and August 1961, respectively. The dark areas correspond to a positive energy balance, that is, an excess of incoming radiation over the outgoing. The light areas indicate regions of negative balance. In these figures, the darkest shade (for example, southwestern United States in July) has a value of $> +1.75 \times 10^5$ ergs $\text{cm}^{-2} \text{sec}^{-1}$, while the lightest shade (for example, 40°N to 50°N belt in December) corresponds to $< -1.65 \times 10^5$ ergs $\text{cm}^{-2} \text{sec}^{-1}$. The intermediate values of the energy balance are plotted, in steps of about 0.5×10^5 ergs $\text{cm}^{-2} \text{sec}^{-1}$, as seven gradations of the shading level.

The two triangular regions comprising parts of South America and Siberia have been left blank because data from Tiros are not available in those regions. Also the several other 10° by 10° grids left blank in March and July are due to nonavailability of data from Tiros for these regions.

A preliminary examination of these figures reveals several interesting features.

1) In December and January, the regions of maximum positive energy input are located in the latitudinal belts of 20°S to 50°S , while in July and August it is the 10°N to 40°N belt which has a high excess of energy. The evolution of this phenomenon is revealed by the charts for February and March.

2) The geographical distribution of the energy balance appears to be such that the desert areas of Africa, Australia, the Middle East, and southwestern United States show extreme positive energy inputs in the local summer.

3) The effect of the monsoon over India is noticeable in comparing the charts for March and August 1961. The net energy input over India is lower in August than in March, presumably because of the heavy monsoon cloud

cover, although if it were not for the monsoon, one would expect a very high excess of energy in the summer month of August.

4) In the Northern Hemisphere, the region of the western Pacific appears to show a relative deficit in energy during all the 6 months. This area is well known for strong cyclogenetic activity. As in the case of the monsoon, the high cloudiness produced by the cyclogenesis probably accounts for the observed low input in energy.

A detailed analysis of these charts, in conjunction with the actually observed global distribution of the weather patterns for the respective months, may provide a better understanding of the role played by the energy balance of the atmosphere in the evolution of weather systems.

S. I. RASOOL

*Goddard Institute for Space Studies,
National Aeronautics and Space
Administration, New York,
and Department of Meteorology,
New York University, New York*

References and Notes

1. G. C. Simpson, *Mem. Roy. Meteorol. Soc.* 3, No. 23, 53 (1929).
2. G. H. Houghton, *J. Meteorol.* 11, (1954).
3. J. London, "A study of the atmospheric heat balance," *Final Report*, Contract No. AF 19 (122)-165, Research Division, College of Engineering, New York University, New York, 1957.
4. F. A. Budyko, Jr., *The Heat Balance of the Earth's Surface*. Translated from the Russian edition by Nina A. Stepanova (U.S. Weather Bureau, Washington, D.C., 1958).
5. W. R. Bandeen, R. A. Hanel, J. Licht, R. A. Stampfl, W. G. Stroud, *J. Geophys. Res.* 65, 3165 (1961).
6. W. Nordberg, W. R. Bandeen, B. J. Conrath, V. Kinde, I. Persano, *J. Atm. Sci.* 19, 20 (1962).
7. A. Arking, *Science*, this issue.
8. D. G. Wark, G. Yamamoto, J. H. Lienesch, "Methods of estimating infrared flux and surface temperature from meteorological satellites," *J. Atmospheric Sci.* 19, 369 (1962).
9. C. Prabhakara and S. I. Rasool, in *Proceedings of the 1st International Symposium on Rocket and Satellite Meteorology*, H. Wexler and J. E. Caskey, Jr., Eds. (North-Holland, Amsterdam, 1963).
10. I thank J. London, R. M. Goody, J. Charney, and R. Jastrow for many illuminating discussions, and Conrad Hipkins who computed and plotted these data on an IBM 7094 and SC 4020.

3 January 1964